


The Nucleus of the Milky Way



ISAAC/VLT 1.3 + 1.71 + 2.09 μm


blue: stars (near-infrared)

green: ionized gas
(radio - cm)

red: molecular gas
(radio - mm)

Circum Nuclear Ring

mini-spiral
(Sagittarius A West)




blue: stars (near-infrared)

green: ionized gas (radio - cm)

red: molecular gas (radio - mm)

Circum Nuclear Ring


mini-spiral
(Sagittarius A West)



blue: stars (near-infrared)

green: ionized gas
(radio - cm)


red: molecular gas
(radio - mm)



blue: stars (near-infrared)

green: ionized gas
(radio - cm)

red: molecular gas
(radio - mm)




ng

blue: stars (near-infrared)

green: ionized gas
(radio - cm)


red: molecular gas
(radio - mm)



blue: stars (near-infrared)

green: ionized gas
(radio - cm)

red: molecular gas
(radio - mm)



50 light days



14 light days



MPE/ESO

UCLA/Keck


e.g. Eckart & Genzel (1996); Ghez et al. (1998, 2003, 2008); Genzel et al. (2000); Eckart et al. (2002); Schödel et al. (2002, 2003); Reid et al. (2004); Eisenhauer et al. (2003, 2005); Gillessen et al. (2009); Yelda et al. (2010)

50 light days



MPE/ESO

14 light days



UCLA/Keck


e.g. Eckart & Genzel (1996); Ghez et al. (1998, 2003, 2008); Genzel et al. (2000); Eckart et al. (2002); Schödel et al. (2002, 2003); Reid et al. (2004); Eisenhauer et al. (2003, 2005); Gillessen et al. (2009); Yelda et al. (2010)

50 light days



MPE/ESO

14 light days



UCLA/Keck

e.g. Eckart & Genzel (1996); Ghez et al. (1998, 2003, 2008); Genzel et al. (2000); Eckart et al. (2002); Schödel et al. (2002, 2003); Reid et al. (2004); Eisenhauer et al. (2003, 2005); Gillessen et al. (2009); Yelda et al. (2010)


50 light days



14 light days




- Distance: 8.1 - 8.3 kpc
 - Mass: $4.1 - 4.3 \times 10^6 M_{\odot}$
 - Mass density: $> 5 \times 10^{15} M_{\odot} \text{ pc}^{-3}$
- \Rightarrow angular size of $R_{\text{Schwarzschild}} \approx 10 \mu\text{as}$
(along with M87 largest BH on the sky)




MPE/ESO

UCLA/Keck

e.g. Eckart & Genzel (1996); Ghez et al. (1998, 2003, 2008); Genzel et al. (2000); Eckart et al. (2002); Schödel et al. (2002, 2003); Reid et al. (2004); Eisenhauer et al. (2003, 2005); Gillessen et al. (2009); Yelda et al. (2010)



NASA/JPL-Caltech/S. Stolovy (Spitzer Science Center/Caltech)




NASA/JPL-Caltech/S. Stolovy (Spitzer Science Center/Caltech)

NSC + central BH:
Is there a stellar
cusp?


Stellar cusps around MBHs

Figure: M. Freitag



Stellar cusps around MBHs


Figure: M. Freitag




2-body relaxed cluster
around massive black
hole should display a
cusp with
 $\rho \propto r^{-1.5 \dots -1.75}$

Stellar cusps around MBHs

Figure: M. Freitag




2-body relaxed cluster
around massive black
hole should display a
cusp with
 $\rho \propto r^{-1.5 \dots -1.75}$



Stellar cusps around MBHs


Figure: M. Freitag



2-body relaxed cluster
around massive black
hole should display a
cusp with
 $\rho \propto r^{-1.5 \dots -1.75}$

Cusp formation is a **robust**
prediction of theoretical stellar
dynamics


(e.g., Lightman & Shapiro, 1977; Bahcall & Wolf
1976, 1977; Freitag+ 2006; Hopman & Alexander
2006).



ISAAC/VLT 1.3 + 2.09 μm
20,000 point sources
(seeing limited, FWHM $\sim 0.4''$)

150" / 18 light years


Sgr A*




ISAAC/VLT 1.3 + 2.09 μm
20,000 point sources
(seeing limited, FWHM $\sim 0.4''$)


150'' / 18 light years

Sgr A*






Stellar surface number density



Schödel et al. (2007)

Stellar surface number density



Schödel et al. (2007)

Classifying stars at the GC

...is painful business.

- high ($A_V \approx 40, A_K \approx 3$) and variable (scales of arcsec) extinction
- only H,K,L observations (narrow range of stellar colors)
- crowded field: ~ 10 sources/arsec²
- FOV of spectroscopy very small
- photometric precision limited by crowding and PSF variability

Classifying stars at the GC


...is painful business.

- high ($A_V \approx 40, A_K \approx 3$) and variable (scales of arcsec) extinction
- only H,K,L observations (narrow range of stellar colors)
- crowded field: ~ 10 sources/arcsec²
- FOV of spectroscopy very small
- photometric precision limited by crowding and PSF variability


... but it can be done

- reasonably accurately with IFS for hundreds of stars (e.g. *Eisenhauer+ 2005; Maness+ 2007; Do+ 2009; Bartko+ 2010*), using dozens of hours of observing time
- more dirtily ($\sim 90\%$ accuracy) but efficiently via narrow band imaging for 1000s of stars in the central pc (*Buchholz + 2009*)

$n(r)$ of old stars \neq $n(r)$ of young stars




$n(r)$ of old stars \neq $n(r)$ of young stars



Stellar classification with AO-assisted integral field spectroscopy

(OSIRIS/Keck Do et al., 2009; see also SINFONI/VLT : Bartko et al. 2010)



Stellar classification with AO-assisted integral field spectroscopy

(OSIRIS/Keck Do et al., 2009; see also SINFONI/VLT : Bartko et al. 2010)

Decreasing density of old stars toward Sgr A.*

→ $\gamma < 1.0$ with $>99\%$ probability *(Do et al., 2009)*

→ There is no observable cusp, maybe even a deficit of old stars around Sgr A*

Known for the brightest ($K < 13$) giants since the 1990s.
The new measurements show that there is also a deficit
of the lower mass RC giants near Sgr A*.

(Sellgren+ 1990; Genzel+ 1996; Haller+ 1996; first indication for RC stars given in Genzel+ 2003)

1


Radius (arcsec)

Where is the stellar cusp at the GC?

- Destroyed:
e.g., by infall of IMBH up to a few 10^9 yr ago (see Merritt & Szell 2006)
- Not yet formed:
necessary time scale may be longer than $\sim 10^{10}$ yr (Merritt 2009)
- Invisible:
giants could be destroyed by collisions with MS stars and BHs in dense cluster center; however, mechanism probably not effective enough (e.g. Dale+ 2009)
- Are our assumptions correct?
Continuous star formation, cluster not old enough?, cluster structure on large scales, relation to nuclear stellar disk, fraction of disrupted stars accreted onto BH?, etc.

Where is the stellar cusp at the GC?

- Destroy
e.g., by in
- Not yet
necessar
- Invisible
giants cc
dense cl
enough (
- Are our
Continu
structur
of disrupt



(Szell 2006)

(t 2009)

and BHs in
: effective

ster
, fraction

Where is the stellar cusp at the GC?

- Destroyed:
e.g., by infall of IMBH up to a few 10^9 yr ago (see Merritt & Szell 2006)
- Not yet formed:
necessary time scale may be longer than $\sim 10^{10}$ yr (Merritt 2009)
- Invisible:
giants could be destroyed by collisions with MS stars and BHs in dense cluster center; however, mechanism probably not effective enough (e.g. Dale+ 2009)
- Are our assumptions correct?
Continuous star formation, cluster not old enough?, cluster structure on large scales, relation to nuclear stellar disk, fraction of disrupted stars accreted onto BH?, etc.

What do we observe at the GC?

50% completeness limit in central
parsec: $K \approx 18$ (MS stars of $\sim 2M_{\odot}$)

Main tracers are HB/RC stars.

What do we observe at the GC?



50% completeness limit in central parsec: $K \approx 18$ (MS stars of $\sim 2M_{\odot}$)

Main tracers are HB/RC stars.

Source: Wikipedia
Created by Uwe Kils (iceberg) and User:Wiska Bodo (sky).